

Product datasheet for RC205651L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Angiopoietin like 4 (ANGPTL4) (NM 139314) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Angiopoietin like 4 (ANGPTL4) (NM_139314) Human Tagged ORF Clone Lentiviral Particle

Symbol: Angiopoietin like 4

Synonyms: ARP4; FIAF; HARP; HFARP; NL2; PGAR; pp1158; TGQTL; UNQ171

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_139314 **ORF Size:** 1218 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC205651).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 139314.1

 RefSeq Size:
 1905 bp

 RefSeq ORF:
 1221 bp

 Locus ID:
 51129

 UniProt ID:
 Q9BY76

 Cytogenetics:
 19p13.2

Protein Families: Druggable Genome, Secreted Protein

Protein Pathways: PPAR signaling pathway





Angiopoietin like 4 (ANGPTL4) (NM_139314) Human Tagged ORF Clone Lentiviral Particle – RC205651L2V

MW: 45.2 kDa

Gene Summary:

This gene encodes a glycosylated, secreted protein containing a C-terminal fibrinogen domain. The encoded protein is induced by peroxisome proliferation activators and functions as a serum hormone that regulates glucose homeostasis, lipid metabolism, and insulin sensitivity. This protein can also act as an apoptosis survival factor for vascular endothelial cells and can prevent metastasis by inhibiting vascular growth and tumor cell invasion. The C-terminal domain may be proteolytically-cleaved from the full-length secreted protein. Decreased expression of this gene has been associated with type 2 diabetes. Alternative splicing results in multiple transcript variants. This gene was previously referred to as ANGPTL2 but has been renamed ANGPTL4. [provided by RefSeq, Sep 2013]