

## Product datasheet for RC205653L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## C1QA (NM\_015991) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** C1QA (NM\_015991) Human Tagged ORF Clone Lentiviral Particle

Symbol: C1QA

Mammalian Cell Puromycin

Selection:

Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

**ACCN:** NM\_015991

ORF Size: 735 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC205653).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 015991.2</u>, <u>NP 057075.1</u>

RefSeq Size: 1098 bp
RefSeq ORF: 738 bp
Locus ID: 712

 UniProt ID:
 P02745

 Cytogenetics:
 1p36.12

**Domains:** C1Q, Collagen **Protein Families:** Secreted Protein

**Protein Pathways:** Complement and coagulation cascades, Prion diseases, Systemic lupus erythematosus





## C1QA (NM\_015991) Human Tagged ORF Clone Lentiviral Particle - RC205653L3V

**MW:** 26 kDa

**Gene Summary:** 

This gene encodes the A-chain polypeptide of serum complement subcomponent C1q, which associates with C1r and C1s to yield the first component of the serum complement system. C1q deficiency is associated with lupus erythematosus and glomerulonephritis. C1q is composed of 18 polypeptide chains which include 6 A-chains, 6 B-chains, and 6 C-chains. Each chain contains an N-terminal collagen-like region and a C-terminal C1q globular domain. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Nov 2016]