

Product datasheet for RC205718L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RRM2 (NM_001034) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: RRM2 (NM_001034) Human Tagged ORF Clone Lentiviral Particle

Symbol: RRM2

Synonyms: C2orf48; R2; RR2; RR2M

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_001034 **ORF Size:** 1167 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC205718).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 001034.1</u>

 RefSeq Size:
 2500 bp

 RefSeq ORF:
 1170 bp

 Locus ID:
 6241

 UniProt ID:
 P31350

 Cytogenetics:
 2p25.1

Domains: ribonuc_red_sm

Protein Families: Druggable Genome





RRM2 (NM_001034) Human Tagged ORF Clone Lentiviral Particle - RC205718L4V

Protein Pathways: Glutathione metabolism, Metabolic pathways, p53 signaling pathway, Purine metabolism,

Pyrimidine metabolism

MW: 44.7 kDa

Gene Summary: This gene encodes one of two non-identical subunits for ribonucleotide reductase. This

reductase catalyzes the formation of deoxyribonucleotides from ribonucleotides. Synthesis of the encoded protein (M2) is regulated in a cell-cycle dependent fashion. Transcription from this gene can initiate from alternative promoters, which results in two isoforms that differ in the lengths of their N-termini. Related pseudogenes have been identified on chromosomes 1

and X. [provided by RefSeq, Sep 2009]