

Product datasheet for RC205723L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Proteasome subunit beta type 4 (PSMB4) (NM_002796) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Proteasome subunit beta type 4 (PSMB4) (NM_002796) Human Tagged ORF Clone Lentiviral

Particle

Symbol: Proteasome subunit beta type 4

Synonyms: HN3; HsN3; PRAAS3; PROS-26; PROS26

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_002796

ORF Size: 792 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC205723).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 002796.2</u>

 RefSeq Size:
 925 bp

 RefSeq ORF:
 795 bp

 Locus ID:
 5692

 UniProt ID:
 P28070

 Cytogenetics:
 1q21.3

Domains: proteasome





Proteasome subunit beta type 4 (PSMB4) (NM_002796) Human Tagged ORF Clone Lentiviral Particle – RC205723L4V

Protein Families: Druggable Genome, Protease

Protein Pathways: Proteasome

MW: 29 kDa

Gene Summary: The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S

core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family,

that is a 20S core beta subunit. [provided by RefSeq, Jul 2008]