

Product datasheet for **RC205727**

Kindlin 2 (FERMT2) (NM_006832) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Kindlin 2 (FERMT2) (NM_006832) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	Kindlin 2
Synonyms:	KIND2; mig-2; MIG2; PLEKHC1; UNC112; UNC112B
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

**ORF Nucleotide
Sequence:**

>RC205727 ORF sequence
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGGCTCTGGACGGGATAAGGATGCCAGATGGCTGCTACGCGGACGGGACGTGGGAACTGAGTGTCCATG
 TGACGGACCTGAACCGCATGTCACCCGAGAGTGACCGCGAGGTGCACATTGGAGGCGTGATGCTTAA
 GCTGGTGGAGAACTCGATGTAAAAAAGATTGGTCTGACCATGCTCTCTGGTGGAAAAGAAGAGAACT
 TGGCTTCTGAAGACACATTGGACCTTAGATAAGTATGGTATTCAGGCAGATGCTAAGCTTCAGTTACCC
 CTCAGCACAACTGCTCCGCTGCAGCTTCCCAACATGAAGTATGTGAAGGTGAAAGTGAATTTCTCTGA
 TAGAGTCTTCAAAGCTGTTTCTGACATCTGTAAGACTTTAATATCAGACACCCCGAAGAAGCTTTCTCTC
 TTAAGAAACCCAGAGATCCAACAAGAAAAAAGAAGAAGCTAGATGACCAGTCTGAAGATGAGGCAC
 TTGAATTAGAGGGCCTTATCACTCCTGGATCAGGAAGTATATATTCAAGCCAGGACTGTATAGTAA
 AACATGACCCCACTTATGATGCTCATGATGGAAGCCCTTGTACCAACTTCTGCTTGGTTTGGTGAC
 AGTGCTTTGTGAGAAGGCAATCCTGGTATACTTGTGTCAGTCAACCAATCACGTACCAGAAATCTTGG
 CAAAAATGTTCAAGCCTCAAGCTCTTCTTGATAAAGCAAAAAATCAACCAAGGATGGCTTGATTCCTCAAG
 ATCTCTCATGGAACAAGATGTGAAGGAAAAATGAGGCCTTGTGCTCCGATTCAAGTATTACAGCTTTTT
 GATTTGAATCCAAAGTATGATGCAATCAGAATCAATCAGCTTTATGAGCAGGCCAAATGGGCCATTTCTC
 TGGAAGAGATTGAATGCACAGAAGAAGAAATGATGATGTTGCAGCCCTGCAGTATCATATCAATAAGCT
 GTCAATCATGACATCAGAGAATCATTGAAACAACAGTGACAAAGAAGTTGATGAAGTTGATGCTGCCCTT
 TCAGACCTGGAGATTACTCTGGAAGGGGTAAAACGTCAACAATTTGGGTGACATTACTCCATTCTCTG
 AACTTGTGACTACATTAAGTTTTCAAGCCAAAAAGCTGACTCTGAAAGGTTACAAACAATATTGGTG
 CACCTTCAAAGACACATCCATTTCTTGTATAAGAGCAAAGAAGAAATCCAGTGGCACACCCAGCTCATCAG
 ATGAACCTCAGGGGATGTGAAGTTACCCAGATGTAAACATTTACGGCCAAAAATTTAACATTAACCTCC
 TGATTCAGTTGCAGAAGGCATGAATGAAATCTGGCTTCGTTGTGACAATGAAAAACAGTATGCACACTG
 GATGGCAGCTGCAGATTAGCCTCAAAGGCAAGACCATGGCGGACAGTTCTTACAACTTAGAAGTTCAG
 AATATTCTTTCTTTCTGAAGATGCAGCATTTAAACCCAGATCCTCAGTTAATACCAGAGCAGATCACGA
 CTGATATAACTCCTGAATGTTTGGTGTCTCCCCGCTATCTAAAAAAGTATAAGAACAAGCAGATAACAGC
 GAGAATCTTGAGGCCCATCAGAATGTAGCTCAGATGAGTCTAATTGAAGCCAAGATGAGATTTATTCAA
 GCTTGGCAGTCACTACCTGAATTTGGCATCACTCACTTATTGCAAGGTTCCAAGGGGGAAAAAAGAAG
 AACTTATTGGAATTGCATACAACAGACTGATTCGGATGGATGCCAGCACTGGAGATGCAATTAACACATG
 GCGTTTCAGCAACATGAAACAGTGAATGTCAACTGGGAAATCAAAATGGTCACCGTAGAGTTTGCAGAT
 GAAGTACGATTGTCTTCATTTGTACTGAAGTAGATTGCAAAGTGGTTTCAATGAAATTCATTTGGTGGCTACA
 TATTTCTCTCAACACGTGCAAAGACCAAACGAGAGTTTAGATGAAGAGATGTTCTACAAACTTACCAG
 TGGTTGGGTG

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Protein Sequence: >RC205727 protein sequence
Red=Cloning site Green=Tags(s)


MALDGIRMPDGCYADGTWELSVHVTDLNRDVTLRVTGEVHIGGVMLKLVKLDVKKDWSHDHALWWEKKRT
 WLLKTHWTLDKYGIQADAKLQFTPOHKLLRLQLPNMKYVKVKNFSDRVFKAUSDICKTFNIRHPEELSL
 LKKPRDPTKKKKKLLDDQSEDEALELEGLITPGSGSIYSSPLYSKTMPTPYDAHDGSPLSPTSAAWFGD
 SALSEGNGILAVSQPITSPEILAKMFKPQALLDKAKINQGWLDSRSLMEQDVKENEALLRKFYYSFF
 DLNPKYDAIRINQLYEQAKWAILLEEIECTEEMMFAALQYHINKLSIMTSENHLNNSDKEVDEVDAAL
 SDLEITLEGGKTSTILGDITSIPELADYIKVFKPKKLTLYGKQYWCTFKDTSISCYKSKEESSGTPAHQ
 MNLRGCEVTPDVNISGQKFNIKLLIPVAEGMNEIWLRCNEKQYAHWMAACRLASKGKTMADSSYNLEVQ
 NILSFLKMQHLPDPQLIPEQITTDITPECLVSPRYLKKYKNKQITARILEAHQNV AQMSLIEAKMRFIQ
 AWQSLPEFGITHFIARFQGGKKEELIGIAYNRLIRMDASTGDAIKTWRFSNMKQVNVNWEIKMVTVEFAD
 EVRLSFICTEVDCKVHEFIGGYIFLSTRAKDQNESLDEEMFYKLTSGWV

TRTRPLEQKLISEEDLANDILDYKDDDDKV

Chromatograms: https://cdn.origene.com/chromatograms/mk6574_d12.zip

Restriction Sites: SgfI-MluI

Cloning Scheme:



ACCN: NM_006832


ORF Size: 2040 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)


OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM_006832.1</u> , <u>NP_006823.1</u>
RefSeq Size:	3351 bp
RefSeq ORF:	2043 bp
Locus ID:	10979
UniProt ID:	<u>Q96AC1</u>
Cytogenetics:	14q22.1
MW:	77.9 kDa
Gene Summary:	Scaffolding protein that enhances integrin activation mediated by TLN1 and/or TLN2, but activates integrins only weakly by itself. Binds to membranes enriched in phosphoinositides. Enhances integrin-mediated cell adhesion onto the extracellular matrix and cell spreading; this requires both its ability to interact with integrins and with phospholipid membranes. Required for the assembly of focal adhesions. Participates in the connection between extracellular matrix adhesion sites and the actin cytoskeleton and also in the orchestration of actin assembly and cell shape modulation. Recruits FBLIM1 to focal adhesions. Plays a role in the TGFB1 and integrin signaling pathways. Stabilizes active CTNNB1 and plays a role in the regulation of transcription mediated by CTNNB1 and TCF7L2/TCF4 and in Wnt signaling. [UniProtKB/Swiss-Prot Function]


Product images:




Circular map for RC205727



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY FERMT2 (Cat# RC205727, Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-FERMT2 (Cat# [TA500504]). Positive lysates [LY416377] (100ug) and [LC416377] (20ug) can be purchased separately from OriGene.



Western blot validation of overexpression lysate (Cat# [LY416377]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC205727 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified FERMT2 protein (Cat# [TP305727]). The protein was produced from HEK293T cells transfected with FERMT2 cDNA clone (Cat# RC205727) using MegaTran 2.0 (Cat# [TT210002]).