

Product datasheet for **RC205753L2V**

STAT5A (NM_003152) Human Tagged ORF Clone Lentiviral Particle

Product data:

| | |
|---------------------------|--|
| Product Type: | Lentiviral Particles |
| Product Name: | STAT5A (NM_003152) Human Tagged ORF Clone Lentiviral Particle |
| Symbol: | STAT5A |
| Synonyms: | MGF; STAT5 |
| Mammalian Cell Selection: | None |
| Vector: | pLenti-C-mGFP (PS100071) |
| Tag: | mGFP |
| ACCN: | NM_003152 |
| ORF Size: | 2382 bp |
| ORF Nucleotide Sequence: | The ORF insert of this clone is exactly the same as(RC205753). |
| OTI Disclaimer: | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info |
| OTI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| RefSeq: | NM_003152.3 , NP_003143.2 |
| RefSeq Size: | 4298 bp |
| RefSeq ORF: | 2385 bp |
| Locus ID: | 6776 |
| UniProt ID: | P42229 |
| Cytogenetics: | 17q21.2 |
| Domains: | SH2, STAT |



[View online »](#)

| | |
|--------------------------|---|
| Protein Families: | Druggable Genome, ES Cell Differentiation/IPS, Stem cell relevant signaling - DSL/Notch pathway, Stem cell relevant signaling - JAK/STAT signaling pathway, Transcription Factors |
| Protein Pathways: | Acute myeloid leukemia, Chronic myeloid leukemia, ErbB signaling pathway, Jak-STAT signaling pathway, Pathways in cancer |
| MW: | 90.5 kDa |
| Gene Summary: | <p>The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated by, and mediates the responses of many cell ligands, such as IL2, IL3, IL7 GM-CSF, erythropoietin, thrombopoietin, and different growth hormones. Activation of this protein in myeloma and lymphoma associated with a TEL/JAK2 gene fusion is independent of cell stimulus and has been shown to be essential for tumorigenesis. The mouse counterpart of this gene is found to induce the expression of BCL2L1/BCL-X(L), which suggests the antiapoptotic function of this gene in cells. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Dec 2013]</p> |