

## Product datasheet for RC205786L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CD32A (FCGR2A) (NM\_021642) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

Product Name: CD32A (FCGR2A) (NM\_021642) Human Tagged ORF Clone Lentiviral Particle

Symbol: CD32A

Synonyms: CD32; CD32A; CDw32; FCG2; FcGR2; FCGR2A1; IGFR2

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_021642

ORF Size: 948 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC205786).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 021642.2

 RefSeq Size:
 2411 bp

 RefSeq ORF:
 951 bp

 Locus ID:
 2212

 UniProt ID:
 P12318

 Cytogenetics:
 1q23.3

 Domains:
 ig, IGc2, IG

Protein Families: ES Cell Differentiation/IPS, Transmembrane





## CD32A (FCGR2A) (NM\_021642) Human Tagged ORF Clone Lentiviral Particle - RC205786L2V

**Protein Pathways:** Fc gamma R-mediated phagocytosis, Systemic lupus erythematosus

**MW:** 34.7 kDa

**Gene Summary:** This gene encodes one member of a family of immunoglobulin Fc receptor genes found on

the surface of many immune response cells. The protein encoded by this gene is a cell surface receptor found on phagocytic cells such as macrophages and neutrophils, and is involved in the process of phagocytosis and clearing of immune complexes. Alternative

splicing results in multiple transcript variants. [provided by RefSeq, Oct 2008]