

## Product datasheet for RC205816L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CTIP1 (BCL11A) (NM 018014) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** CTIP1 (BCL11A) (NM\_018014) Human Tagged ORF Clone Lentiviral Particle

Symbol:

BCL11A-L; BCL11a-M; BCL11A-S; BCL11A-XL; CTIP1; DILOS; EVI9; HBFQTL5; ZNF856 Synonyms:

**Mammalian Cell** 

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK NM 018014 ACCN: **ORF Size:** 2319 bp

**ORF Nucleotide** 

Sequence:

OTI Disclaimer:

**Domains:** 

The ORF insert of this clone is exactly the same as(RC205816).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 018014.2, NP 060484.2

zf-C2H2

RefSeq Size: 3958 bp RefSeq ORF: 2322 bp Locus ID: 53335 **UniProt ID:** Q9H165 Cytogenetics: 2p16.1

**Protein Families: Transcription Factors** 





ORIGENE

**MW:** 83.9 kDa

Gene Summary:

This gene encodes a C2H2 type zinc-finger protein by its similarity to the mouse Bcl11a/Evi9 protein. The corresponding mouse gene is a common site of retroviral integration in myeloid leukemia, and may function as a leukemia disease gene, in part, through its interaction with BCL6. During hematopoietic cell differentiation, this gene is down-regulated. It is possibly involved in lymphoma pathogenesis since translocations associated with B-cell malignancies also deregulates its expression. Multiple transcript variants encoding several different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]