

## Product datasheet for RC205818L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## CHST8 (NM\_022467) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** CHST8 (NM\_022467) Human Tagged ORF Clone Lentiviral Particle

Symbol: CHST8

**Synonyms:** GalNAc4ST; GALNAC4ST1; PSS3

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM\_022467

ORF Size: 1272 bp

**ORF Nucleotide** 

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(RC205818).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 022467.3, NP 071912.2

 RefSeq Size:
 2496 bp

 RefSeq ORF:
 1275 bp

 Locus ID:
 64377

 UniProt ID:
 Q9H2A9

 Cytogenetics:
 19q13.11

**Domains:** Sulfotransfer2

**Protein Families:** Transmembrane





ORIGENE

MW: 48.8 kDa

**Gene Summary:** The protein encoded by this gene belongs to the sulfotransferase 2 family. It is

predominantly expressed in the pituitary gland, and is localized to the golgi membrane. This protein catalyzes the transfer of sulfate to position 4 of non-reducing N-acetylgalactosamine (GalNAc) residues in both N-glycans and O-glycans. It is responsible for sulfation of GalNAc on luteinizing hormone (LH), which is required for production of the sex hormones. Mice lacking this enzyme, exhibit increased levels of circulating LH, and precocious sexual maturation of both male and female mice. Alternatively spliced transcript variants have been

found for this gene. [provided by RefSeq, Aug 2011]