

## Product datasheet for **RC205983L3V**

### NSF (NM\_006178) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

|                           |  |
|---------------------------|--|
| Product Type:             | Lentiviral Particles   |
| Product Name:             | NSF (NM_006178) Human Tagged ORF Clone Lentiviral Particle   |
| Symbol:                   | NSF  |
| Synonyms:                 | SEC18; SKD2  |
| Mammalian Cell Selection: | Puromycin  |
| Vector:                   | pLenti-C-Myc-DDK-P2A-Puro (PS100092)   |
| Tag:                      | Myc-DDK  |
| ACCN:                     | NM_006178  |
| ORF Size:                 | 2232 bp  |
| ORF Nucleotide Sequence:  | The ORF insert of this clone is exactly the same as(RC205983).   |
| OTI Disclaimer:           | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a> |
| OTI Annotation:           | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.   |
| RefSeq:                   | <a href="#">NM_006178.1</a>  |
| RefSeq Size:              | 3995 bp  |
| RefSeq ORF:               | 2235 bp  |
| Locus ID:                 | 4905   |
| UniProt ID:               | <a href="#">P46459</a>   |
| Cytogenetics:             | 17q21.31   |
| Domains:                  | cdc48_N, AAA, AAA, cdc48_2   |
| Protein Families:         | Protease   |



[View online »](#)

**MW:** 82.6 kDa

**Gene Summary:** Required for vesicle-mediated transport. Catalyzes the fusion of transport vesicles within the Golgi cisternae. Is also required for transport from the endoplasmic reticulum to the Golgi stack. Seems to function as a fusion protein required for the delivery of cargo proteins to all compartments of the Golgi stack independent of vesicle origin. Interaction with AMPAR subunit GRIA2 leads to influence GRIA2 membrane cycling (By similarity).[UniProtKB/Swiss-Prot Function]