

## Product datasheet for RC206057L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## NPY1R (NM\_000909) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** NPY1R (NM\_000909) Human Tagged ORF Clone Lentiviral Particle

Symbol: NPY1R

Synonyms: NPY1-R; NPYR

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

NM 000909

Tag: Myc-DDK

ORF Size: 1152 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC206057).

Sequence:

ACCN:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 000909.4</u>

 RefSeq Size:
 2974 bp

 RefSeq ORF:
 1155 bp

 Locus ID:
 4886

 UniProt ID:
 P25929

Cytogenetics: 4q32.2

**Domains:** 7tm 1

**Protein Families:** Druggable Genome, GPCR, Transmembrane





## NPY1R (NM\_000909) Human Tagged ORF Clone Lentiviral Particle - RC206057L3V

**Protein Pathways:** Neuroactive ligand-receptor interaction

MW: 44.4 kDa

**Gene Summary:** This gene belongs to the G-protein-coupled receptor superfamily. The encoded

transmembrane protein mediates the function of neuropeptide Y (NPY), a neurotransmitter, and peptide YY (PYY), a gastrointestinal hormone. The encoded receptor undergoes fast agonist-induced internalization through clathrin-coated pits and is subsequently recycled back to the cell membrane. Activation of Y1 receptors may result in mobilization of intracellular calcium and inhibition of adenylate cyclase activity. [provided by RefSeq, Aug

2013]