

Product datasheet for RC206132L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

GYLTL1B (LARGE2) (NM 152312) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: GYLTL1B (LARGE2) (NM_152312) Human Tagged ORF Clone Lentiviral Particle

Symbol:

GYLTL1B; PP5656 Synonyms:

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

mGFP Tag:

NM 152312 ACCN: **ORF Size:** 2163 bp

OTI Disclaimer:

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC206132).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 152312.3

RefSeq Size: 2539 bp RefSeq ORF: 2166 bp Locus ID: 120071 **UniProt ID:** Q8N3Y3 Cytogenetics: 11p11.2

Protein Families: Transmembrane

MW: 81.6 kDa







Gene Summary:

Bifunctional glycosyltransferase with both xylosyltransferase and beta-1,3-glucuronyltransferase activities involved in the biosynthesis of the phosphorylated O-mannosyl trisaccharide (N-acetylgalactosamine-beta-3-N-acetylglucosamine-beta-4-(phosphate-6-)mannose), a carbohydrate structure present in alpha-dystroglycan (DAG1). Phosphorylated O-mannosyl trisaccharid is required for binding laminin G-like domain-containing extracellular proteins with high affinity. Elongates the glucuronyl-beta-1,4-xylose-beta disaccharide primer structure by adding repeating units [-3-Xylose-alpha-1,3-GlcA-beta-1-] to produce a heteropolysaccharide. Has a higher activity toward alpha-dystroglycan than LARGE.[UniProtKB/Swiss-Prot Function]