

#### OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for RC206235L3

### Silencer of Death Domain (BAG4) (NM\_004874) Human Tagged Lenti ORF Clone

#### **Product data:**

| Product Type:                | Expression Plasmids  |
|------------------------------|--|
| Product Name:                | Silencer of Death Domain (BAG4) (NM_004874) Human Tagged Lenti ORF Clone                         |
| Tag:                         | Myc-DDK  |
| Symbol:                      | Silencer of Death Domain   |
| Synonyms:                    | BAG-4; SODD  |
| Mammalian Cell<br>Selection: | Puromycin  |
| Vector:                      | pLenti-C-Myc-DDK-P2A-Puro (PS100092)   |
| E. coli Selection:           | Chloramphenicol (34 ug/mL)   |
| ORF Nucleotide<br>Sequence:  | The ORF insert of this clone is exactly the same as(RC206235).                                   |
| <b>Restriction Sites:</b>    | Sgfl-Mlul  |
| Cloning Scheme:              |  |
|                              | Cloning sites used for ORF Shuttling:  |
|                              | Sgf I         ORF         Mlu I            GCG ATC GCC         ATG//         NNN         ACG CGT |



ACCN: ORF Size: NM\_004874 1371 bp




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

|                        | er of Death Domain (BAG4) (NM_004874) Human Tagged Lenti ORF Clone – RC206235L3   |
|------------------------|---|
| OTI Disclaimer:        | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>   |
| OTI Annotation:        | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.  |
| Components:            | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).  |
| Reconstitution Method: | <ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>  |
| RefSeq:                | <u>NM 004874.2</u>  |
| RefSeq Size:           | 4478 bp   |
| RefSeq ORF:            | 1374 bp   |
| Locus ID:              | 9530  |
| UniProt ID:            | <u>095429</u>   |
| Cytogenetics:          | 8p11.23   |
| Domains:               | BAG   |
| Protein Families:      | Druggable Genome  |
| MW:                    | 49.6 kDa  |
| Gene Summary:          | The protein encoded by this gene is a member of the BAG1-related protein family. BAG1 is an anti-apoptotic protein that functions through interactions with a variety of cell apoptosis and growth related proteins including BCL-2, Raf-protein kinase, steroid hormone receptors, growth factor receptors and members of the heat shock protein 70 kDa family. This protein contains a BAG domain near the C-terminus, which could bind and inhibit the chaperone activity of Hsc70/Hsp70. This protein was found to be associated with the death domain of tumor necrosis factor receptor type 1 (TNF-R1) and death receptor-3 (DR3), and thereby negatively regulates downstream cell death signaling. The regulatory role of this protein in cell death was demonstrated in epithelial cells which undergo apoptosis while integrin mediated matrix contacts are lost. Alternatively spliced transcript variants encoding distinct |

isoforms have been identified. [provided by RefSeq, Mar 2011]

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

## **Product images:**



Circular map for RC206235L3

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US