

## Product datasheet for RC206271L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Heparin Cofactor II (SERPIND1) (NM 000185) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** Heparin Cofactor II (SERPIND1) (NM\_000185) Human Tagged ORF Clone Lentiviral Particle

Symbol: Heparin Cofactor II

D22S673; HC2; HCF2; HCII; HLS2; LS2; THPH10 Synonyms:

**Mammalian Cell** 

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK NM 000185 ACCN:

**ORF Size:** 1497 bp

**ORF Nucleotide** 

OTI Disclaimer:

Sequence:

The ORF insert of this clone is exactly the same as(RC206271).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 000185.3

RefSeq Size: 2237 bp RefSeq ORF: 1500 bp Locus ID: 3053 **UniProt ID:** P05546 Cytogenetics: 22q11.21

**Domains: SERPIN** 

**Protein Families:** Druggable Genome





## Heparin Cofactor II (SERPIND1) (NM\_000185) Human Tagged ORF Clone Lentiviral Particle – RC206271L3V

**Protein Pathways:** Complement and coagulation cascades

**MW:** 57 kDa

**Gene Summary:** This gene belongs to the serpin gene superfamily. Serpins play roles in many processes

including inflammation, blood clotting, and cancer metastasis. Members of this family have highly conserved secondary structures with a reactive center loop that interacts with the protease active site to inhibit protease activity. This gene encodes a plasma serine protease that functions as a thrombin and chymotrypsin inhibitor. The protein is activated by heparin, dermatan sulfate, and glycosaminoglycans. Allelic variations in this gene are associated with

heparin cofactor II deficiency. [provided by RefSeq, Jul 2015]