

Product datasheet for **RC206300**

GABPA (NM_002040) Human Tagged ORF Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	GABPA (NM_002040) Human Tagged ORF Clone
Tag:	Myc-DDK
Symbol:	GABPA
Synonyms:	E4TF1-60; E4TF1A; NFT2; NRF2; NRF2A; RCH04A07
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Cloning Scheme:


ACCN: NM_002040

ORF Size: 1362 bp

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

- Reconstitution Method:**
1. Centrifuge at 5,000xg for 5min.
 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
 3. Close the tube and incubate for 10 minutes at room temperature.
 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_002040.4](#)

RefSeq Size: 5182 bp

RefSeq ORF: 1365 bp

Locus ID: 2551

UniProt ID: [Q06546](#)

Cytogenetics: 21q21.3


Domains: ETS, SAM_PNT

Protein Families: Transcription Factors


MW: 51.3 kDa

Gene Summary: This gene encodes one of three GA-binding protein transcription factor subunits which functions as a DNA-binding subunit. Since this subunit shares identity with a subunit encoding the nuclear respiratory factor 2 gene, it is likely involved in activation of cytochrome oxidase expression and nuclear control of mitochondrial function. This subunit also shares identity with a subunit constituting the transcription factor E4TF1, responsible for expression of the adenovirus E4 gene. Because of its chromosomal localization and ability to form heterodimers with other polypeptides, this gene may play a role in the Down Syndrome phenotype. Two transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Oct 2010]


Product images:



Circular map for RC206300



Western blot validation of overexpression lysate (Cat# [LY400747]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC206300 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).



Coomassie blue staining of purified GABPA protein (Cat# [TP306300]). The protein was produced from HEK293T cells transfected with GABPA cDNA clone (Cat# RC206300) using MegaTran 2.0 (Cat# [TT210002]).