

Product datasheet for RC206379L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Caspase 10 (CASP10) (NM_032977) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Caspase 10 (CASP10) (NM_032977) Human Tagged ORF Clone Lentiviral Particle

Symbol: CASP10

Synonyms: ALPS2; FLICE-2; FLICE2; MCH4

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_032977 **ORF Size:** 1566 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC206379).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 032977.3

RefSeq Size: 5906 bp
RefSeq ORF: 1569 bp
Locus ID: 843

 UniProt ID:
 Q92851

 Cytogenetics:
 2q33.1

Domains: Peptidase_C14, DED

Protein Families: Druggable Genome, Protease





Protein Pathways: Apoptosis, RIG-I-like receptor signaling pathway

MW: 59 kDa

Gene Summary: This gene encodes a protein which is a member of the cysteine-aspartic acid protease

(caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein cleaves and activates caspases 3 and 7, and the protein itself is processed by caspase 8. Mutations in this gene are associated with type

IIA autoimmune lymphoproliferative syndrome, non-Hodgkin lymphoma and gastric cancer. Alternatively spliced transcript variants encoding different isoforms have been described for

this gene. [provided by RefSeq, Apr 2011]