

Product datasheet for RC206420L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IL3RA (NM_002183) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: IL3RA (NM_002183) Human Tagged ORF Clone Lentiviral Particle

Symbol: IL3RA

Synonyms: CD123; hlL-3Ra; lL3RAY; lL3RX; lL3RX

Mammalian Cell

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK
ACCN: NM_002183

ORF Size: 1134 bp

ORF Nucleotide

OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC206420).

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 002183.2</u>

 RefSeq Size:
 1726 bp

 RefSeq ORF:
 1137 bp

 Locus ID:
 3563

 UniProt ID:
 P26951

Cytogenetics: X;Y

Protein Families: Transmembrane





IL3RA (NM_002183) Human Tagged ORF Clone Lentiviral Particle - RC206420L1V

Protein Pathways: Apoptosis, Cytokine-cytokine receptor interaction, Hematopoietic cell lineage, Jak-STAT

signaling pathway

MW: 43.33 kDa

Gene Summary: The protein encoded by this gene is an interleukin 3 specific subunit of a heterodimeric

cytokine receptor. The receptor is comprised of a ligand specific alpha subunit and a signal transducing beta subunit shared by the receptors for interleukin 3 (IL3), colony stimulating factor 2 (CSF2/GM-CSF), and interleukin 5 (IL5). The binding of this protein to IL3 depends on the beta subunit. The beta subunit is activated by the ligand binding, and is required for the biological activities of IL3. This gene and the gene encoding the colony stimulating factor 2

receptor alpha chain (CSF2RA) form a cytokine receptor gene cluster in a X-Y

pseudoautosomal region on chromosomes X or Y. Alternatively spliced transcript variants

encoding distinct isoforms have been found. [provided by RefSeq, Jun 2012]