

## Product datasheet for RC206420L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## IL3RA (NM 002183) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type: Lentiviral Particles** 

**Product Name:** IL3RA (NM\_002183) Human Tagged ORF Clone Lentiviral Particle

Symbol:

CD123; hIL-3Ra; IL3R; IL3RAY; IL3RX; IL3RY Synonyms:

**Mammalian Cell** 

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

mGFP Tag:

NM 002183 ACCN: **ORF Size:** 1134 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC206420).

Sequence: OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 002183.2

RefSeq Size: 1726 bp RefSeq ORF: 1137 bp Locus ID: 3563 **UniProt ID:** P26951 X:Y

Cytogenetics:

**Protein Families:** Transmembrane





## IL3RA (NM\_002183) Human Tagged ORF Clone Lentiviral Particle - RC206420L2V

**Protein Pathways:** Apoptosis, Cytokine-cytokine receptor interaction, Hematopoietic cell lineage, Jak-STAT

signaling pathway

MW: 43.33 kDa

**Gene Summary:** The protein encoded by this gene is an interleukin 3 specific subunit of a heterodimeric

cytokine receptor. The receptor is comprised of a ligand specific alpha subunit and a signal transducing beta subunit shared by the receptors for interleukin 3 (IL3), colony stimulating factor 2 (CSF2/GM-CSF), and interleukin 5 (IL5). The binding of this protein to IL3 depends on the beta subunit. The beta subunit is activated by the ligand binding, and is required for the biological activities of IL3. This gene and the gene encoding the colony stimulating factor 2

receptor alpha chain (CSF2RA) form a cytokine receptor gene cluster in a X-Y

pseudoautosomal region on chromosomes X or Y. Alternatively spliced transcript variants

encoding distinct isoforms have been found. [provided by RefSeq, Jun 2012]