

## Product datasheet for RC206450L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## WFDC1 (NM 021197) Human Tagged ORF Clone Lentiviral Particle

## **Product data:**

**Product Type: Lentiviral Particles** 

**Product Name:** WFDC1 (NM 021197) Human Tagged ORF Clone Lentiviral Particle

Symbol: PS20 Synonyms: **Mammalian Cell** 

Selection:

None

pLenti-C-Myc-DDK (PS100064) Vector:

Myc-DDK Tag: NM 021197 ACCN:

**ORF Size:** 660 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC206450).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 021197.2

RefSeq Size: 1396 bp RefSeq ORF: 663 bp Locus ID: 58189 **UniProt ID:** Q9HC57 Cytogenetics: 16q24.1 **Domains:** WAP

**Protein Families:** Secreted Protein





ORIGENE

**MW:** 20.6 kDa

**Gene Summary:** 

This gene encodes a member of the WAP-type four disulfide core domain family. The WAP-type four-disulfide core domain contains eight cysteines forming four disulfide bonds at the core of the protein, and functions as a protease inhibitor in many family members. This gene is mapped to chromosome 16q24, an area of frequent loss of heterozygosity in cancers, including prostate, breast and hepatocellular cancers and Wilms' tumor. This gene is downregulated in many cancer types and may be involved in the inhibition of cell proliferation. The encoded protein may also play a role in the susceptibility of certain CD4 memory T cells to human immunodeficiency virus infection. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2013]