

Product datasheet for RC206450L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

WFDC1 (NM_021197) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: WFDC1 (NM_021197) Human Tagged ORF Clone Lentiviral Particle

Symbol: WFDC1
Synonyms: PS20
Mammalian Cell None

- ·

Selection:

INOTIC

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_021197

ORF Size: 660 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC206450).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 021197.2

RefSeq Size: 1396 bp
RefSeq ORF: 663 bp
Locus ID: 58189
UniProt ID: Q9HC57
Cytogenetics: 16q24.1
Domains: WAP

Protein Families: Secreted Protein





ORIGENE

MW: 20.6 kDa

Gene Summary:

This gene encodes a member of the WAP-type four disulfide core domain family. The WAP-type four-disulfide core domain contains eight cysteines forming four disulfide bonds at the core of the protein, and functions as a protease inhibitor in many family members. This gene is mapped to chromosome 16q24, an area of frequent loss of heterozygosity in cancers, including prostate, breast and hepatocellular cancers and Wilms' tumor. This gene is downregulated in many cancer types and may be involved in the inhibition of cell proliferation. The encoded protein may also play a role in the susceptibility of certain CD4 memory T cells to human immunodeficiency virus infection. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2013]