

Product datasheet for RC206464L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Factor XIIIa (F13A1) (NM_000129) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Factor XIIIa (F13A1) (NM_000129) Human Tagged ORF Clone Lentiviral Particle

Symbol: Factor XIIIa

Synonyms: F13A

Mammalian Cell None

Selection:

Vector: pLenti-C-Myc-DDK (PS100064)

Tag: Myc-DDK

ACCN: NM_000129 **ORF Size:** 2196 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC206464).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000129.3

 RefSeq Size:
 3863 bp

 RefSeq ORF:
 2199 bp

 Locus ID:
 2162

 UniProt ID:
 P00488

Cytogenetics: 6p25.1

Domains: Transglutamin_C, TGc

Protein Families: Druggable Genome, Secreted Protein



defective wound healing, and habitual abortion. [provided by RefSeq, Jul 2008]



Protein Pathways: Complement and coagulation cascades

MW: 83.3 kDa

Gene Summary: This gene encodes the coagulation factor XIII A subunit. Coagulation factor XIII is the last

zymogen to become activated in the blood coagulation cascade. Plasma factor XIII is a heterotetramer composed of 2 A subunits and 2 B subunits. The A subunits have catalytic function, and the B subunits do not have enzymatic activity and may serve as plasma carrier molecules. Platelet factor XIII is comprised only of 2 A subunits, which are identical to those of plasma origin. Upon cleavage of the activation peptide by thrombin and in the presence of calcium ion, the plasma factor XIII dissociates its B subunits and yields the same active enzyme, factor XIIIa, as platelet factor XIII. This enzyme acts as a transglutaminase to catalyze the formation of gamma-glutamyl-epsilon-lysine crosslinking between fibrin molecules, thus stabilizing the fibrin clot. It also crosslinks alpha-2-plasmin inhibitor, or fibronectin, to the alpha chains of fibrin. Factor XIII deficiency is classified into two categories: type I deficiency, characterized by the lack of both the A and B subunits; and type II deficiency, characterized by the lack of the A subunit alone. These defects can result in a lifelong bleeding tendency,