

Product datasheet for RC206523L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Macrophage Inflammatory Protein 3 beta (CCL19) (NM_006274) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Macrophage Inflammatory Protein 3 beta (CCL19) (NM_006274) Human Tagged ORF Clone

Lentiviral Particle

Symbol: Macrophage Inflammatory Protein 3 beta

Synonyms: CKb11; ELC; MIP-3b; MIP3B; SCYA19

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM_006274

ORF Size: 294 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC206523).

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 006274.2

 RefSeq Size:
 684 bp

 RefSeq ORF:
 297 bp

 Locus ID:
 6363

 UniProt ID:
 Q99731

 Cytogenetics:
 9p13.3

Protein Families: Druggable Genome, Secreted Protein





Macrophage Inflammatory Protein 3 beta (CCL19) (NM_006274) Human Tagged ORF Clone Lentiviral Particle – RC206523L3V

Protein Pathways: Chemokine signaling pathway, Cytokine-cytokine receptor interaction

MW: 11 kDa

Gene Summary: This antimicrobial gene is one of several CC cytokine genes clustered on the p-arm of

chromosome 9. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent

cysteines. The cytokine encoded by this gene may play a role in normal lymphocyte recirculation and homing. It also plays an important role in trafficking of T cells in thymus, and in T cell and B cell migration to secondary lymphoid organs. It specifically binds to

chemokine receptor CCR7. [provided by RefSeq, Sep 2014]