

## Product datasheet for RC206531L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## RAMP2 (NM\_005854) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** RAMP2 (NM\_005854) Human Tagged ORF Clone Lentiviral Particle

Symbol: RAMP2

Mammalian Cell Puromycin

Selection:

Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK

ACCN: NM\_005854

ORF Size: 525 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as (RC206531).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 005854.2</u>

RefSeq Size: 808 bp

RefSeq ORF: 528 bp

Locus ID: 10266

UniProt ID: 060895

Cytogenetics: 17q21.2

Domains: RAMP

Protein Families: Druggable Genome, Transmembrane

**Protein Pathways:** Vascular smooth muscle contraction





ORIGENE

**MW:** 19.6 kDa

**Gene Summary:** 

The protein encoded by this gene is a member of the RAMP family of single-transmembrane-domain proteins, called receptor (calcitonin) activity modifying proteins (RAMPs). RAMPs are type I transmembrane proteins with an extracellular N terminus and a cytoplasmic C terminus. RAMPs are required to transport calcitonin-receptor-like receptor (CRLR) to the plasma membrane. CRLR, a receptor with seven transmembrane domains, can function as either a calcitonin-gene-related peptide (CGRP) receptor or an adrenomedullin receptor, depending on which members of the RAMP family are expressed. In the presence of this (RAMP2) protein, CRLR functions as an adrenomedullin receptor. The RAMP2 protein is involved in core glycosylation and transportation of adrenomedullin receptor to the cell surface. [provided by RefSeq, Jul 2008]