

## Product datasheet for RC206536L3V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## P2X6 (P2RX6) (NM\_005446) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** P2X6 (P2RX6) (NM\_005446) Human Tagged ORF Clone Lentiviral Particle

Symbol: P2X6

Synonyms: P2RXL1; P2X6; P2XM

Mammalian Cell

Selection:

Puromycin

**Vector:** pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM\_005446

ORF Size: 1323 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC206536).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 005446.3</u>

 RefSeq Size:
 2754 bp

 RefSeq ORF:
 1326 bp

 Locus ID:
 9127


 UniProt ID:
 015547

 Cytogenetics:
 22q11.21

**Protein Families:** Druggable Genome, Ion Channels: ATP Receptors, Transmembrane

**Protein Pathways:** Calcium signaling pathway, Neuroactive ligand-receptor interaction





**ORÏGENE** 

MW: 48.6 kDa

**Gene Summary:** The protein encoded by this gene belongs to the family of P2X receptors, which are ATP-gated

ion channels and mediate rapid and selective permeability to cations. This gene is

predominantly expressed in skeletal muscle, and regulated by p53. The encoded protein is associated with VE-cadherin at the adherens junctions of human umbilical vein endothelial cells. Alternative splicing results in multiple transcript variants. A related pseudogene, which is also located on chromosome 22, has been identified. [provided by RefSeq, Apr 2009]