

Product datasheet for RC206545L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

HABP2 (NM_004132) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: HABP2 (NM 004132) Human Tagged ORF Clone Lentiviral Particle

Symbol: HABP2

Synonyms: FSAP; HABP; HGFAL; NMTC5; PHBP

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_004132 **ORF Size:** 1680 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC206545).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 004132.2</u>

 RefSeq Size:
 3019 bp

 RefSeq ORF:
 1683 bp

 Locus ID:
 3026

 UniProt ID:
 Q14520

 Cytogenetics:
 10q25.3

Domains: KR, Tryp_SPc, EGF, EGF

Protein Families: Druggable Genome, Protease, Secreted Protein





ORIGENE

MW: 62.7 kDa

Gene Summary:

This gene encodes a member of the peptidase S1 family of serine proteases. The encoded preproprotein is secreted by hepatocytes and proteolytically processed to generate heavy and light chains that form the mature heterodimer. Further autoproteolysis leads to smaller, inactive peptides. This extracellular protease binds hyaluronic acid and may play a role in the coagulation and fibrinolysis systems. Mutations in this gene are associated with nonmedullary thyroid cancer and susceptibility to venous thromboembolism. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed. [provided by RefSeq, Jan 2016]