

Product datasheet for RC206589L1

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Nociceptin receptor (OPRL1) (NM_182647) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: Nociceptin receptor (OPRL1) (NM_182647) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: Nociceptin receptor

Synonyms: KOR-3; KOR3; NOCIR; NOP; NOPr; OOR; OPRL; ORL1; PNOCR

Mammalian Cell None

Selection:

Vector:pLenti-C-Myc-DDK (PS100064)E. coli Selection:Chloramphenicol (34 ug/mL)

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC206589).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_182647

ORF Size: 1110 bp



Nociceptin receptor (OPRL1) (NM_182647) Human Tagged Lenti ORF Clone - RC206589L1

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 182647.1</u>

 RefSeq Size:
 3391 bp

 RefSeq ORF:
 1113 bp

 Locus ID:
 4987

 UniProt ID:
 P41146

 Cytogenetics:
 20q13.33

Protein Families: Druggable Genome, GPCR, Transmembrane

Protein Pathways: Neuroactive ligand-receptor interaction

MW: 40.7 kDa


Gene Summary: The protein encoded by this gene is a member of the 7 transmembrane-spanning G protein-

coupled receptor family, and functions as a receptor for the endogenous, opioid-related neuropeptide, nociceptin/orphanin FQ. This receptor-ligand system modulates a variety of biological functions and neurobehavior, including stress responses and anxiety behavior, learning and memory, locomotor activity, and inflammatory and immune responses. A promoter region between this gene and the 5'-adjacent RGS19 (regulator of G-protein signaling 19) gene on the opposite strand functions bi-directionally as a core-promoter for both genes, suggesting co-operative transcriptional regulation of these two functionally related genes. Alternatively spliced transcript variants have been described for this gene. A recent study provided evidence for translational readthrough in this gene, and expression of an additional C-terminally extended isoform via the use of an alternative in-frame translation


termination codon. [provided by RefSeq, Dec 2017]



Product images:



Circular map for RC206589L1



Double digestion of RC206589L1 using Sgfl and Mlul $\,$