

# Product datasheet for RC206596L1V

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# CXCR3 (NM\_001504) Human Tagged ORF Clone Lentiviral Particle

#### **Product data:**

**Product Type:** Lentiviral Particles

**Product Name:** CXCR3 (NM\_001504) Human Tagged ORF Clone Lentiviral Particle

Symbol: CXCR3

Synonyms: CD182; CD183; CKR-L2; CMKAR3; GPR9; IP10-R; Mig-R; MigR

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

Tag:Myc-DDKACCN:NM\_001504

ORF Size: 1104 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC206596).

Sequence:
OTI Disclaimer:

**Domains:** 

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 001504.1

 RefSeq Size:
 1670 bp

 RefSeq ORF:
 1107 bp

 Locus ID:
 2833

 UniProt ID:
 P49682

 Cytogenetics:
 Xq13.1

Protein Families: Druggable Genome, GPCR, Transmembrane

7tm 1





## CXCR3 (NM\_001504) Human Tagged ORF Clone Lentiviral Particle - RC206596L1V

**Protein Pathways:** Chemokine signaling pathway, Cytokine-cytokine receptor interaction

**MW:** 40.5 kDa

**Gene Summary:** This gene encodes a G protein-coupled receptor with selectivity for three chemokines, termed

CXCL9/Mig (monokine induced by interferon-g), CXCL10/IP10 (interferon-g-inducible 10 kDa protein) and CXCL11/I-TAC (interferon-inducible T cell a-chemoattractant). Binding of chemokines to this protein induces cellular responses that are involved in leukocyte traffic,

most notably integrin activation, cytoskeletal changes and chemotactic migration.

Alternatively spliced transcript variants encoding different isoforms have been found for this gene. One of the isoforms (CXCR3-B) shows high affinity binding to chemokine, CXCL4/PF4

(PMID:12782716). [provided by RefSeq, Jun 2011]