

Product datasheet for RC206756L1V

1062 bp

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ARG2 (NM_001172) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: ARG2 (NM_001172) Human Tagged ORF Clone Lentiviral Particle

Symbol: ARG2 **Mammalian Cell** None

Selection:

Vector:

pLenti-C-Myc-DDK (PS100064)

Myc-DDK Tag: ACCN: NM_001172 **ORF Size:**

ORF Nucleotide

Sequence:

OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC206756).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

This clone was engineered to express the complete ORF with an expression tag. Expression **OTI Annotation:**

varies depending on the nature of the gene.

RefSeq: NM 001172.3

RefSeq Size: 1981 bp RefSeq ORF: 1065 bp Locus ID: 384 **UniProt ID:** P78540

Cytogenetics: 14q24.1 Domains: arginase

Protein Pathways: Arginine and proline metabolism, Metabolic pathways

MW: 38.6 kDa







Gene Summary:

Arginase catalyzes the hydrolysis of arginine to ornithine and urea. At least two isoforms of mammalian arginase exists (types I and II) which differ in their tissue distribution, subcellular localization, immunologic crossreactivity and physiologic function. The type II isoform encoded by this gene, is located in the mitochondria and expressed in extra-hepatic tissues, especially kidney. The physiologic role of this isoform is poorly understood; it is thought to play a role in nitric oxide and polyamine metabolism. Transcript variants of the type II gene resulting from the use of alternative polyadenylation sites have been described. [provided by RefSeq, Jul 2008]