

## Product datasheet for RC206777L1V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## HEXO (ERI1) (NM\_153332) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** HEXO (ERI1) (NM\_153332) Human Tagged ORF Clone Lentiviral Particle

Symbol: HEXC

Synonyms: 3'HEXO; HEXO; THEX1

Mammalian Cell

Selection:

None

**Vector:** pLenti-C-Myc-DDK (PS100064)

 Tag:
 Myc-DDK

 ACCN:
 NM\_153332

 ORF Size:
 1047 bp

**ORF Nucleotide** 

OTI Disclaimer:

1047 bp

Sequence:

The ORF insert of this clone is exactly the same as(RC206777).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 153332.2

 RefSeq Size:
 4615 bp

 RefSeq ORF:
 1050 bp

 Locus ID:
 90459

 UniProt ID:
 Q8IV48

 Cytogenetics:
 8p23.1

**MW:** 40.1 kDa







## **Gene Summary:**

RNA exonuclease that binds to the 3'-end of histone mRNAs and degrades them, suggesting that it plays an essential role in histone mRNA decay after replication. A 2' and 3'-hydroxyl groups at the last nucleotide of the histone 3'-end is required for efficient degradation of RNA substrates. Also able to degrade the 3'-overhangs of short interfering RNAs (siRNAs) in vitro, suggesting a possible role as regulator of RNA interference (RNAi). Requires for binding the 5'-ACCCA-3' sequence present in stem-loop structure. Able to bind other mRNAs. Required for 5.8S rRNA 3'-end processing. Also binds to 5.8s ribosomal RNA. Binds with high affinity to the stem-loop structure of replication-dependent histone pre-mRNAs.[UniProtKB/Swiss-Prot Function]