

Product datasheet for RC206920L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Activin A Receptor Type IB (ACVR1B) (NM_004302) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Activin A Receptor Type IB (ACVR1B) (NM_004302) Human Tagged ORF Clone Lentiviral

Particle

Symbol: Activin A Receptor Type IB

Synonyms: ACTRIB; ACVRLK4; ALK4; SKR2

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_004302

ORF Size: 1515 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as (RC206920).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 004302.3</u>

RefSeq Size: 4540 bp RefSeq ORF: 1518 bp

Locus ID: 91

 UniProt ID:
 P36896

 Cytogenetics:
 12q13.13

Domains: Activin_recp, pkinase, TyrKc, S_TKc, GS





Activin A Receptor Type IB (ACVR1B) (NM_004302) Human Tagged ORF Clone Lentiviral Particle – RC206920L2V

Protein Families: Druggable Genome, Protein Kinase, Transmembrane

Protein Pathways: Adherens junction, Chronic myeloid leukemia, Colorectal cancer, Cytokine-cytokine receptor

interaction, Endocytosis, MAPK signaling pathway, Pancreatic cancer, Pathways in cancer,

TGF-beta signaling pathway

MW: 56.81 kDa

Gene Summary: This gene encodes an activin A type IB receptor. Activins are dimeric growth and

differentiation factors which belong to the transforming growth factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I and two type II receptors. This protein is a type I receptor which is essential for signaling. Mutations in this gene are associated with pituitary tumors. Alternate splicing results in multiple transcript variants.

[provided by RefSeq, Jun 2010]