

## Product datasheet for RC206974L2V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## TORC1 (CRTC1) (NM\_015321) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

**Product Type:** Lentiviral Particles

**Product Name:** TORC1 (CRTC1) (NM\_015321) Human Tagged ORF Clone Lentiviral Particle

Symbol: TORC1

Synonyms: Mam-2; MAML2; MECT1; TORC-1; TORC1; WAMTP1

**Mammalian Cell** 

Selection:

None

**Vector:** pLenti-C-mGFP (PS100071)

Tag: mGFP

**ACCN:** NM\_015321 **ORF Size:** 1902 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC206974).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeg:** NM 015321.1, NP 056136.1

 RefSeq Size:
 2505 bp

 RefSeq ORF:
 1905 bp

 Locus ID:
 23373

 UniProt ID:
 Q6UUV9

 Cytogenetics:
 19p13.11

**Protein Families:** Transcription Factors

**MW:** 67.1 kDa





## **Gene Summary:**

Transcriptional coactivator for CREB1 which activates transcription through both consensus and variant cAMP response element (CRE) sites. Acts as a coactivator, in the SIK/TORC signaling pathway, being active when dephosphorylated and acts independently of CREB1 'Ser-133' phosphorylation. Enhances the interaction of CREB1 with TAF4. Regulates the expression of specific CREB-activated genes such as the steroidogenic gene, StAR. Potent coactivator of PGC1alpha and inducer of mitochondrial biogenesis in muscle cells. In the hippocampus, involved in late-phase long-term potentiation (L-LTP) maintenance at the Schaffer collateral-CA1 synapses. May be required for dendritic growth of developing cortical neurons (By similarity). In concert with SIK1, regulates the light-induced entrainment of the circadian clock. In response to light stimulus, coactivates the CREB-mediated transcription of PER1 which plays an important role in the photic entrainment of the circadian clock. [UniProtKB/Swiss-Prot Function]