

Product datasheet for RC207025L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Beta TRCP (BTRC) (NM_033637) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Beta TRCP (BTRC) (NM_033637) Human Tagged ORF Clone Lentiviral Particle

Symbol: Beta TRCP

Synonyms: BETA-TRCP; betaTrCP; bTrCP1; FBW1A; FBXW1; FBXW1A; FWD1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 033637

ORF Size: 1815 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC207025).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 033637.2

 RefSeq Size:
 6180 bp

 RefSeq ORF:
 1818 bp

 Locus ID:
 8945

 UniProt ID:
 Q9Y297

 Cytogenetics:
 10q24.32

Domains: WD40, F-box

Protein Families: Druggable Genome





Beta TRCP (BTRC) (NM_033637) Human Tagged ORF Clone Lentiviral Particle - RC207025L3V

Protein Pathways: Hedgehog signaling pathway, Oocyte meiosis, Ubiquitin mediated proteolysis, Wnt signaling

pathway

MW: 68.9 kDa

Gene Summary: This gene encodes a member of the F-box protein family which is characterized by an

approximately 40 amino acid motif, the F-box. The F-box proteins constitute one of the four subunits of ubiquitin protein ligase complex called SCFs (SKP1-cullin-F-box), which function in phosphorylation-dependent ubiquitination. The F-box proteins are divided into 3 classes: Fbws containing WD-40 domains, Fbls containing leucine-rich repeats, and Fbxs containing either different protein-protein interaction modules or no recognizable motifs. The protein encoded by this gene belongs to the Fbws class; in addition to an F-box, this protein contains multiple WD-40 repeats. The encoded protein mediates degradation of CD4 via its interaction with HIV-1 Vpu. It has also been shown to ubiquitinate phosphorylated NFKBIA (nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha), targeting it for degradation and thus activating nuclear factor kappa-B. Alternatively spliced transcript variants have been described. A related pseudogene exists in chromosome 6. [provided by RefSeq, Mar 2012]