

Product datasheet for RC207113L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Thymine DNA glycosylase (TDG) (NM_003211) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Thymine DNA glycosylase (TDG) (NM 003211) Human Tagged ORF Clone Lentiviral Particle

Symbol: Thymine DNA glycosylase

Synonyms: hTDG

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_003211

 ORF Size:
 1230 bp

ORF Nucleotide

OTI Disclaimer:

1230 bp

Sequence:

Domains:

The ORF insert of this clone is exactly the same as(RC207113).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 003211.3

 RefSeq Size:
 3251 bp

 RefSeq ORF:
 1233 bp

 Locus ID:
 6996

 UniProt ID:
 Q13569

 Cytogenetics:
 12q23.3

Protein Families: Druggable Genome

UDG





Thymine DNA glycosylase (TDG) (NM_003211) Human Tagged ORF Clone Lentiviral Particle – RC207113L3V

Protein Pathways: Base excision repair

MW: 46 kDa

Gene Summary: The protein encoded by this gene belongs to the TDG/mug DNA glycosylase family. Thymine-

DNA glycosylase (TDG) removes thymine moieties from G/T mismatches by hydrolyzing the carbon-nitrogen bond between the sugar-phosphate backbone of DNA and the mispaired

thymine. With lower activity, this enzyme also removes thymine from C/T and T/T

mispairings. TDG can also remove uracil and 5-bromouracil from mispairings with guanine. This enzyme plays a central role in cellular defense against genetic mutation caused by the

spontaneous deamination of 5-methylcytosine and cytosine. This gene may have a

pseudogene in the p arm of chromosome 12. [provided by RefSeq, Jul 2008]