

## Product datasheet for **RC207117L1V**

### MNDA (NM\_002432) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	MNDA (NM_002432) Human Tagged ORF Clone Lentiviral Particle
Symbol:	MNDA
Synonyms:	PYHIN3
Mammalian Cell Selection:	None
Vector:	pLenti-C-Myc-DDK (PS100064)
Tag:	Myc-DDK
ACCN:	NM_002432
ORF Size:	1221 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC207117).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_002432.1</a>
RefSeq Size:	1670 bp
RefSeq ORF:	1224 bp
Locus ID:	4332
UniProt ID:	<a href="#">P41218</a>
Cytogenetics:	1q23.1
Domains:	PAAD_DAPIN, HIN
Protein Families:	Transcription Factors



[View online »](#)

**MW:** 45.8 kDa

**Gene Summary:** The myeloid cell nuclear differentiation antigen (MNDA) is detected only in nuclei of cells of the granulocyte-monocyte lineage. A 200-amino acid region of human MNDA is strikingly similar to a region in the proteins encoded by a family of interferon-inducible mouse genes, designated Ifi-201, Ifi-202, and Ifi-203, that are not regulated in a cell- or tissue-specific fashion. The 1.8-kb MNDA mRNA, which contains an interferon-stimulated response element in the 5-prime untranslated region, was significantly upregulated in human monocytes exposed to interferon alpha. MNDA is located within 2,200 kb of FCER1A, APCS, CRP, and SPTA1. In its pattern of expression and/or regulation, MNDA resembles IFI16, suggesting that these genes participate in blood cell-specific responses to interferons. [provided by RefSeq, Jul 2008]