

Product datasheet for RC207192L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

FIG4 (NM_014845) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: FIG4 (NM_014845) Human Tagged ORF Clone Lentiviral Particle

Symbol: FIG4

Synonyms: ALS11; BTOP; CMT4J; dJ249I4.1; KIAA0274; SAC3; YVS

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_014845 **ORF Size:** 2721 bp

ORF Nucleotide

OTI Disclaimer:

_. _.

Sequence:

The ORF insert of this clone is exactly the same as(RC207192).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 014845.4</u>

RefSeq Size: 3123 bp RefSeq ORF: 2724 bp

Locus ID: 9896

UniProt ID: Q92562

Cytogenetics: 6q21

Domains: Syja_N

MW: 103.6 kDa







Gene Summary:

The protein encoded by this gene belongs to the SAC domain-containing protein gene family. The SAC domain, approximately 400 amino acids in length and consisting of seven conserved motifs, has been shown to possess phosphoinositide phosphatase activity. The yeast homolog, Sac1p, is involved in the regulation of various phosphoinositides, and affects diverse cellular functions such as actin cytoskeleton organization, Golgi function, and maintenance of vacuole morphology. Membrane-bound phosphoinositides function as signaling molecules and play a key role in vesicle trafficking in eukaryotic cells. Mutations in this gene have been associated with Charcot-Marie-Tooth disease, type 4J. [provided by RefSeq, Jul 2008]