

Product datasheet for **RC207223L3V**

KCNA3 (NM_002232) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type:	Lentiviral Particles
Product Name:	KCNA3 (NM_002232) Human Tagged ORF Clone Lentiviral Particle
Symbol:	KCNA3
Synonyms:	HGK5; HLK3; HPCN3; HUKIII; KV1.3; MK3; PCN3
Mammalian Cell Selection:	Puromycin
Vector:	pLenti-C-Myc-DDK-P2A-Puro (PS100092)
Tag:	Myc-DDK
ACCN:	NM_002232
ORF Size:	1725 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC207223).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	NM_002232.3
RefSeq Size:	3346 bp
RefSeq ORF:	1728 bp
Locus ID:	3738
UniProt ID:	P22001
Cytogenetics:	1p13.3
Domains:	BTB, K_tetra, ion_trans
Protein Families:	Druggable Genome, Ion Channels: Potassium, Transmembrane


[View online »](#)

MW: 63.7 kDa

Gene Summary: Potassium channels represent the most complex class of voltage-gated ion channels from both functional and structural standpoints. Their diverse functions include regulating neurotransmitter release, heart rate, insulin secretion, neuronal excitability, epithelial electrolyte transport, smooth muscle contraction, and cell volume. Four sequence-related potassium channel genes - shaker, shaw, shab, and shal - have been identified in *Drosophila*, and each has been shown to have human homolog(s). This gene encodes a member of the potassium channel, voltage-gated, shaker-related subfamily. This member contains six membrane-spanning domains with a shaker-type repeat in the fourth segment. It belongs to the delayed rectifier class, members of which allow nerve cells to efficiently repolarize following an action potential. It plays an essential role in T-cell proliferation and activation. This gene appears to be intronless and it is clustered together with KCNA2 and KCNA10 genes on chromosome 1. [provided by RefSeq, Jul 2008]