

Product datasheet for RC207230L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CACNB1 (NM_000723) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: CACNB1 (NM_000723) Human Tagged ORF Clone Lentiviral Particle

Symbol: CACNB1

Synonyms: CAB1; CACNLB1; CCHLB1

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_000723 **ORF Size:** 1794 bp

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC207230).

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000723.3

RefSeq Size:3687 bpRefSeq ORF:1797 bp

 Locus ID:
 782

 UniProt ID:
 Q02641

Cytogenetics: 17q12

Domains: Ca_channel_B, SH3, GuKc

Protein Families: Druggable Genome, Ion Channels: Other





CACNB1 (NM_000723) Human Tagged ORF Clone Lentiviral Particle - RC207230L2V

Protein Pathways: Arrhythmogenic right ventricular cardiomyopathy (ARVC), Cardiac muscle contraction, Dilated

cardiomyopathy, Hypertrophic cardiomyopathy (HCM), MAPK signaling pathway

MW: 65.5 kDa

Gene Summary: The protein encoded by this gene belongs to the calcium channel beta subunit family. It plays

an important role in the calcium channel by modulating G protein inhibition, increasing peak calcium current, controlling the alpha-1 subunit membrane targeting and shifting the voltage dependence of activation and inactivation. Alternative splicing occurs at this locus and three transcript variants encoding three distinct isoforms have been identified. [provided by

RefSeq, Jul 2008]