

Product datasheet for RC207294L2V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MRCL3 (MYL12A) (NM 006471) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: MRCL3 (MYL12A) (NM_006471) Human Tagged ORF Clone Lentiviral Particle

Symbol: MRCL3

Synonyms: HEL-S-24; MLC-2B; MLCB; MRCL3; MRLC3; MYL2B

Mammalian Cell

Selection:

None

Vector: pLenti-C-mGFP (PS100071)

Tag: mGFP

ACCN: NM_006471

ORF Size: 513 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC207294).

Sequence:

OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 006471.2, NP 006462.1

 RefSeq Size:
 1243 bp

 RefSeq ORF:
 516 bp

 Locus ID:
 10627

 UniProt ID:
 P19105

 Cytogenetics:
 18p11.31

Domains: EFh





MRCL3 (MYL12A) (NM_006471) Human Tagged ORF Clone Lentiviral Particle - RC207294L2V

Protein Pathways: Focal adhesion, Leukocyte transendothelial migration, Regulation of actin cytoskeleton, Tight

junction

MW: 19.8 kDa

Gene Summary: This gene encodes a nonsarcomeric myosin regulatory light chain. This protein is activated by

phosphorylation and regulates smooth muscle and non-muscle cell contraction. This protein may also be involved in DNA damage repair by sequestering the transcriptional regulator apoptosis-antagonizing transcription factor (AATF)/Che-1 which functions as a repressor of p53-driven apoptosis. Alternate splicing results in multiple transcript variants. A pseudogene

of this gene is found on chromosome 8.[provided by RefSeq, Dec 2014]