

Product datasheet for RC207313

PLAC2 (TINCR) (NM_153375) Human Tagged ORF Clone

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| Product Type: | Expression Plasmids |
|------------------------------|---|
| Product Name: | PLAC2 (TINCR) (NM_153375) Human Tagged ORF Clone |
| Tag: | Myc-DDK |
| Symbol: | PLAC2 |
| Synonyms: | FLJ90734 |
| Mammalian Cell Selection: | Neomycin |
| Vector: | pCMV6-Entry (PS100001) |
| E. coli Selection: | Kanamycin (25 ug/mL) |
| ORF Nucleotide Sequence: | <pre>>RC207313 representing NM_153375 Red=Cloning site Blue=ORF Green=Tags(s)</pre> |
| | TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C |
| | ATGAAACCCTGGCCTGACCAGCTGATGCACACTGCTTCAGACACTCCTGCTGGAGCCCCAGTCCCTGACA AGGACCTAGGACATTTTTGCTCCTGCCCAGCCTATCGGGAGGGA |
| | ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT ACAAGGATGACGACGATAAGG TTTAA |
| Protein Sequence: | >RC207313 representing NM_153375 Red=Cloning site Green=Tags(s) |
| | MKPWPDQLMHTASDTPAGAPVPDKDLGHFCSCPAYREGALSLSALLCDFEVVAPLLGPWVPCLQWKALCH LGKLPWAARGHRKKRSTEGAGETLRGTSALAQDILPATPLPKPCPLPGALFNLFSAAPIHLGLQSPPGSL LRFPPSQPEST |
| | TRTRPLEQKLISEEDLAANDILDYKDDDDKV |
| Chromatograms: | https://cdn.origene.com/chromatograms/mk6546_a04.zip |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


PLAC2 (TINCR) (NM_153375) Human Tagged ORF Clone – RC207313

| Restriction Sites: | Sgfl-Mlul |
|------------------------|--|
| Cloning Scheme: | Cloning sites used for ORF Shuttling: Sgf1 ORF Miul GCGATCGC C MTG NNN ACG CGT |
| | $\frac{Kozac}{Consensus}$ CTATAGGGCGGCGGGAATTCGTCGACTGGATCGGGTACCGGAGGGGATTCGCCGCGGATCGC C DITG |
| | EcoR V Flag.Tag Pme I Fse I GAT CTG GCA GCA AAT GAT ATC CTG GAT TAC AAG GAT GAC GAC GAT AAG GTT TAA ACGGCCGGGCC D L A A N D I L D Y K D D D K V stop |
| | * The last codon before the Stop codon of the ORF |
| CCN: | NM_153375 |
| PRF Size: | 453 bp |
| TI Disclaimer: | The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u> |
| TI Annotation: | This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene. |
| omponents: | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water). |
| Reconstitution Method: | Centrifuge at 5,000xg for 5min. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. Close the tube and incubate for 10 minutes at room temperature. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. |
| RefSeq: | NM 153375.1, NP 699206.1 |
| efSeq Size: | 2467 bp |
| efSeq ORF: | 455 bp |
| ocus ID: | 257000 |
| | |
| | |
| | |
| | |
| | |


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

| | PLAC2 (TINCR) (NM_153375) Human Tagged ORF Clone – RC207313 |
|---------------|--|
| Cytogenetics: | 19p13.3 |
| MW: | 16.5 kDa |
| Gene Summary: | This gene produces a spliced long non-coding RNA that binds RNAs. This transcript interacts with staufen-1 protein to regulate the stability of mRNAs for genes involved in the differentiation of epidermal tissue. Variation in this gene may be associated with cancer progression. [provided by RefSeq, Aug 2017] |

Product images:




Circular map for RC207313



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY PLAC2 (Cat# RC207313, Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PLAC2 (Cat# [TA810461])(1:2000). Positive lysates [LY407086] (100ug) and [LC407086] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Western blot validation of overexpression lysate (Cat# [LY407086]) using anti-DDK antibody (Cat# [TA50011-100]). Left: Cell lysates from untransfected HEK293T cells; Right: Cell lysates from HEK293T cells transfected with RC207313 using transfection reagent MegaTran 2.0 (Cat# [TT210002]).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US