

Product datasheet for RC207315L1

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SIRP alpha (SIRPA) (NM_080792) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: SIRP alpha (SIRPA) (NM_080792) Human Tagged Lenti ORF Clone

Tag: Myc-DDK
Symbol: SIRP alpha

Synonyms: BIT; CD172A; MFR; MYD-1; P84; PTPNS1; SHPS1; SIRP

Mammalian Cell None

Selection:

Vector:pLenti-C-Myc-DDK (PS100064)E. coli Selection:Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC207315).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





^{*} The last codon before the Stop codon of the ORF

ACCN: NM_080792

ORF Size: 1509 bp



SIRP alpha (SIRPA) (NM_080792) Human Tagged Lenti ORF Clone - RC207315L1

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 080792.1

 RefSeq Size:
 3868 bp

 RefSeq ORF:
 1515 bp

 Locus ID:
 140885

 UniProt ID:
 P78324

 Cytogenetics:
 20p13

Domains: ig, IGv, IGc1, IG

Protein Families: Druggable Genome, Phosphatase, Transmembrane


MW: 54.8 kDa

Gene Summary: The protein encoded by this gene is a member of the signal-regulatory-protein (SIRP) family,


and also belongs to the immunoglobulin superfamily. SIRP family members are receptor-type transmembrane glycoproteins known to be involved in the negative regulation of receptor tyrosine kinase-coupled signaling processes. This protein can be phosphorylated by tyrosine kinases. The phospho-tyrosine residues of this PTP have been shown to recruit SH2 domain containing tyrosine phosphatases (PTP), and serve as substrates of PTPs. This protein was found to participate in signal transduction mediated by various growth factor receptors. CD47 has been demonstrated to be a ligand for this receptor protein. This gene and its product share very high similarity with several other members of the SIRP family. These related genes are located in close proximity to each other on chromosome 20p13. Multiple alternatively spliced transcript variants have been determined for this gene. [provided by RefSeq, Jul 2008]



Product images:



Circular map for RC207315L1



Double digestion of RC207315L1 using Sgfl and Mlul