

Product datasheet for RC207328L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SFRP1 (NM_003012) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: SFRP1 (NM_003012) Human Tagged ORF Clone Lentiviral Particle

Symbol: SFRP1

Synonyms: FRP; FRP-1; FRP1; FrzA; SARP2

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_003012

ORF Size: 942 bp

ORF Nucleotide

OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC207328).

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 003012.3</u>

 RefSeq Size:
 4465 bp

 RefSeq ORF:
 945 bp

 Locus ID:
 6422

 UniProt ID:
 Q8N474

 Cytogenetics:
 8p11.21

Domains: FRI, NTR





SFRP1 (NM_003012) Human Tagged ORF Clone Lentiviral Particle - RC207328L4V

Protein Families: Adult stem cells, Cancer stem cells, Druggable Genome, ES Cell Differentiation/IPS, Secreted

Protein, Stem cell relevant signaling - Wnt Signaling pathway, Transmembrane

Protein Pathways: Wnt signaling pathway

MW: 35.2 kDa

Gene Summary: This gene encodes a member of the SFRP family that contains a cysteine-rich domain

homologous to the putative Wnt-binding site of Frizzled proteins. Members of this family act

as soluble modulators of Wnt signaling; epigenetic silencing of SFRP genes leads to

deregulated activation of the Wnt-pathway which is associated with cancer. This gene may also be involved in determining the polarity of photoreceptor cells in the retina. [provided by

RefSeq, Sep 2009]