

Product datasheet for RC207333L3

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

EIF4E (NM_001968) Human Tagged Lenti ORF Clone

Product data:

Product Type: Expression Plasmids

Product Name: EIF4E (NM_001968) Human Tagged Lenti ORF Clone

Tag: Myc-DDK

Symbol: EIF4E

Synonyms: AUTS19; CBP; eIF-4E; EIF4E1; EIF4EL1; EIF4F

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

E. coli Selection: Chloramphenicol (34 ug/mL)

ORF Nucleotide The ORF insert of this clone is exactly the same as(RC207333).

Sequence:

Restriction Sites: Sgfl-Mlul

Cloning Scheme:





 $[\]ensuremath{^*}$ The last codon before the Stop codon of the ORF.

ACCN: NM_001968

ORF Size: 651 bp





EIF4E (NM_001968) Human Tagged Lenti ORF Clone - RC207333L3

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001968.2</u>

RefSeq Size: 4749 bp

RefSeg ORF: 654 bp

Locus ID: 1977

UniProt ID: P06730

Cytogenetics: 4q23

Domains: IF4E

Protein Pathways: Insulin signaling pathway, mTOR signaling pathway

MW: 25.1 kDa


Gene Summary: The protein encoded by this gene is a component of the eukaryotic translation initiation

factor 4F complex, which recognizes the 7-methylguanosine cap structure at the 5' end of messenger RNAs. The encoded protein aids in translation initiation by recruiting ribosomes to the 5'-cap structure. Association of this protein with the 4F complex is the rate-limiting step in translation initiation. This gene acts as a proto-oncogene, and its expression and activation is associated with transformation and tumorigenesis. Several pseudogenes of this gene are found on other chromosomes. Alternative splicing results in multiple transcript variants.

[provided by RefSeq, Sep 2015]



Product images:



Circular map for RC207333L3