

Product datasheet for RC207417L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cytochrome p450 2J2 (CYP2J2) (NM_000775) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: Cytochrome p450 2J2 (CYP2J2) (NM_000775) Human Tagged ORF Clone Lentiviral Particle

Symbol: Cytochrome p450 2J2

Synonyms: CPJ2; CYPIIJ2

Mammalian Cell Puromycin

Selection:

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_000775 **ORF Size:** 1506 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC207417).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 000775.2

 RefSeq Size:
 1876 bp

 RefSeq ORF:
 1509 bp

 Locus ID:
 1573

 UniProt ID:
 P51589

 Cytogenetics:
 1p32.1

Domains: p450

Protein Families: Druggable Genome, P450, Transmembrane





Cytochrome p450 2J2 (CYP2J2) (NM_000775) Human Tagged ORF Clone Lentiviral Particle – RC207417L4V

Protein Pathways: Arachidonic acid metabolism, Linoleic acid metabolism, Metabolic pathways

MW: 57.6 kDa

Gene Summary: This gene encodes a member of the cytochrome P450 superfamily of enzymes. The

cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and is thought to be the predominant enzyme responsible for epoxidation of endogenous arachidonic acid in cardiac tissue. Multiple transcript variants

have been found for this gene. [provided by RefSeq, Jan 2016]