

Product datasheet for RC207481L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

C20orf77 (RPRD1B) (NM 021215) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: C20orf77 (RPRD1B) (NM_021215) Human Tagged ORF Clone Lentiviral Particle

Symbol: C20orf77

Synonyms: C20orf77; CREPT; dJ1057B20.2; K-H; Kub5-Hera; NET60

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

Tag: Myc-DDK
ACCN: NM 021215

ORF Size: 978 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC207481).

OTI Disclaimer:

Sequence:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: <u>NM 021215.2</u>

 RefSeq Size:
 3895 bp

 RefSeq ORF:
 981 bp

 Locus ID:
 58490

 UniProt ID:
 Q9NQG5

 Cytogenetics:
 20q11.23

 Domains:
 RPR, DUF618

MW: 36.9 kDa





Gene Summary:

Interacts with phosphorylated C-terminal heptapeptide repeat domain (CTD) of the largest RNA polymerase II subunit POLR2A, and participates in dephosphorylation of the CTD by RPAP2. Transcriptional regulator which enhances expression of CCND1. Promotes binding of RNA polymerase II to the CCDN1 promoter and to the termination region before the poly-A site but decreases its binding after the poly-A site. Prevents RNA polymerase II from reading through the 3' end termination site and may allow it to be recruited back to the promoter through promotion of the formation of a chromatin loop. Also enhances the transcription of a number of other cell cycle-related genes including CDK2, CDK4, CDK6 and cyclin-E but not CDKN1A, CDKN1B or cyclin-A. Promotes cell proliferation.[UniProtKB/Swiss-Prot Function]