

## Product datasheet for RC207539L4V

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Apc2 (ANAPC2) (NM\_013366) Human Tagged ORF Clone Lentiviral Particle

**Product data:** 

Product Type: Lentiviral Particles

**Product Name:** Apc2 (ANAPC2) (NM\_013366) Human Tagged ORF Clone Lentiviral Particle

Symbol: Apc2 Synonyms: APC2

Mammalian Cell F

Selection:

Puromycin

**Vector:** pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

**ACCN:** NM\_013366 **ORF Size:** 2466 bp

**ORF Nucleotide** 

The ORF insert of this clone is exactly the same as(RC207539).

Sequence:
OTI Disclaimer:

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

**OTI Annotation:** This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

**RefSeq:** <u>NM 013366.3</u>

RefSeq Size: 2733 bp
RefSeq ORF: 2469 bp
Locus ID: 29882
UniProt ID: Q9UJX6
Cytogenetics: 9q34.3
Domains: CULLIN

**Protein Families:** Druggable Genome





## Apc2 (ANAPC2) (NM\_013366) Human Tagged ORF Clone Lentiviral Particle - RC207539L4V

Protein Pathways: Cell cycle, Oocyte meiosis, Progesterone-mediated oocyte maturation, Ubiquitin mediated

proteolysis

**MW:** 93.8 kDa

**Gene Summary:** A large protein complex, termed the anaphase-promoting complex (APC), or the cyclosome,

promotes metaphase-anaphase transition by ubiquitinating its specific substrates such as mitotic cyclins and anaphase inhibitor, which are subsequently degraded by the 26S proteasome. Biochemical studies have shown that the vertebrate APC contains eight subunits. The composition of the APC is highly conserved in organisms from yeast to humans. The product of this gene is a component of the complex and shares sequence similarity with a recently identified family of proteins called cullins, which may also be

involved in ubiquitin-mediated degradation. [provided by RefSeq, Jul 2008]