

Product datasheet for RC207595L1V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TULP3 (NM 003324) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: TULP3 (NM_003324) Human Tagged ORF Clone Lentiviral Particle

Symbol: TUBL3 Synonyms: **Mammalian Cell**

Selection:

None

Vector: pLenti-C-Myc-DDK (PS100064)

Myc-DDK Tag: NM 003324 ACCN: **ORF Size:** 1326 bp

OTI Disclaimer:

ORF Nucleotide

Sequence:

The ORF insert of this clone is exactly the same as(RC207595).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeq: NM 003324.3

RefSeq Size: 3106 bp RefSeq ORF: 1329 bp Locus ID: 7289 <u>0753</u>86 **UniProt ID:** Cytogenetics: 12p13.33

Domains: Tub

Protein Families: Druggable Genome, Transcription Factors





ORIGENE

MW: 49.7 kDa

Gene Summary:

This gene encodes a member of the tubby gene family of bipartite transcription factors. Members of this family have been identified in plants, vertebrates, and invertebrates, and they share a conserved N-terminal transcription activation region and a conserved C-terminal DNA and phosphatidylinositol-phosphate binding region. The encoded protein binds to phosphoinositides in the plasma membrane via its C-terminal region and probably functions as a membrane-bound transcription regulator that translocates to the nucleus in response to phosphoinositide hydrolysis, for instance, induced by G-protein-coupled-receptor signaling. It plays an important role in neuronal development and function. Two transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq, May 2009]