

Product datasheet for RC207595L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TULP3 (NM_003324) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: TULP3 (NM 003324) Human Tagged ORF Clone Lentiviral Particle

Symbol: TULP3
Synonyms: TUBL3

Mammalian Cell Puromycin

Selection: Vector:

pLenti-C-Myc-DDK-P2A-Puro (PS100092)

 Tag:
 Myc-DDK

 ACCN:
 NM_003324

 ORF Size:
 1326 bp

ORF Nucleotide

Sequence:

OTI Disclaimer:

The ORF insert of this clone is exactly the same as(RC207595).

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through

naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 003324.3

 RefSeq Size:
 3106 bp

 RefSeq ORF:
 1329 bp

 Locus ID:
 7289

 UniProt ID:
 075386

 Cytogenetics:
 12p13.33

Domains: Tub

Protein Families: Druggable Genome, Transcription Factors





ORÏGENE

MW: 49.7 kDa

Gene Summary:

This gene encodes a member of the tubby gene family of bipartite transcription factors. Members of this family have been identified in plants, vertebrates, and invertebrates, and they share a conserved N-terminal transcription activation region and a conserved C-terminal DNA and phosphatidylinositol-phosphate binding region. The encoded protein binds to phosphoinositides in the plasma membrane via its C-terminal region and probably functions as a membrane-bound transcription regulator that translocates to the nucleus in response to phosphoinositide hydrolysis, for instance, induced by G-protein-coupled-receptor signaling. It plays an important role in neuronal development and function. Two transcript variants encoding distinct isoforms have been identified for this gene. [provided by RefSeq, May 2009]