

## Product datasheet for **RC207596L2V**

### TRAIL (TNFSF10) (NM\_003810) Human Tagged ORF Clone Lentiviral Particle

#### Product data:

Product Type:	Lentiviral Particles
Product Name:	TRAIL (TNFSF10) (NM_003810) Human Tagged ORF Clone Lentiviral Particle
Symbol:	TRAIL
Synonyms:	Apo-2L; APO2L; CD253; TL2; TNLG6A; TRAIL
Mammalian Cell Selection:	None
Vector:	pLenti-C-mGFP (PS100071)
Tag:	mGFP
ACCN:	NM_003810
ORF Size:	843 bp
ORF Nucleotide Sequence:	The ORF insert of this clone is exactly the same as(RC207596).
OTI Disclaimer:	The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <a href="#">More info</a>
OTI Annotation:	This clone was engineered to express the complete ORF with an expression tag. Expression varies depending on the nature of the gene.
RefSeq:	<a href="#">NM_003810.2</a> , <a href="#">NP_003801.1</a>
RefSeq Size:	1953 bp
RefSeq ORF:	846 bp
Locus ID:	8743
UniProt ID:	<a href="#">P50591</a>
Cytogenetics:	3q26.31
Domains:	TNF
Protein Families:	Druggable Genome, Transmembrane



[View online »](#)

<b>Protein Pathways:</b>	Apoptosis, Cytokine-cytokine receptor interaction, Natural killer cell mediated cytotoxicity
<b>MW:</b>	32.5 kDa
<b>Gene Summary:</b>	<p>The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This protein preferentially induces apoptosis in transformed and tumor cells, but does not appear to kill normal cells although it is expressed at a significant level in most normal tissues. This protein binds to several members of TNF receptor superfamily including TNFRSF10A/TRAILR1, TNFRSF10B/TRAILR2, TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4, and possibly also to TNFRSF11B/OPG. The activity of this protein may be modulated by binding to the decoy receptors TNFRSF10C/TRAILR3, TNFRSF10D/TRAILR4, and TNFRSF11B/OPG that cannot induce apoptosis. The binding of this protein to its receptors has been shown to trigger the activation of MAPK8/JNK, caspase 8, and caspase 3. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2010]</p>