

Product datasheet for RC207597L4V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TLR2 (NM_003264) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: TLR2 (NM_003264) Human Tagged ORF Clone Lentiviral Particle

Symbol: TLR2

Synonyms: CD282; TIL4

Mammalian Cell Puromycin

Selection:

Fulbiliyelli

Vector: pLenti-C-mGFP-P2A-Puro (PS100093)

Tag: mGFP

ACCN: NM_003264 **ORF Size:** 2352 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC207597).

Sequence:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 003264.3

 RefSeq Size:
 3417 bp

 RefSeq ORF:
 2355 bp

 Locus ID:
 7097

 UniProt ID:
 060603

 Cytogenetics:
 4q31.3

Domains: TIR, LRRCT, LRR, LRR_TYP, LRR_BAC, LRR_PS

Protein Families: Druggable Genome, Transmembrane





TLR2 (NM_003264) Human Tagged ORF Clone Lentiviral Particle - RC207597L4V

Protein Pathways: Toll-like receptor signaling pathway

MW: 89.8 kDa

Gene Summary: The protein encoded by this gene is a member of the Toll-like receptor (TLR) family which

plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. This protein is a cell-surface protein that can form heterodimers with other TLR family members to recognize conserved molecules derived from microorganisms known as pathogen-associated molecular patterns (PAMPs). Activation of TLRs by PAMPs leads to an up-regulation of signaling pathways to modulate the host's inflammatory response. This gene is also thought to promote apoptosis in response to bacterial lipoproteins. This gene has been implicated in the pathogenesis of several autoimmune diseases. Alternative splicing

results in multiple transcript variants. [provided by RefSeq, Jan 2016]