

Product datasheet for RC207646L3V

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TAB3 (NM_152787) Human Tagged ORF Clone Lentiviral Particle

Product data:

Product Type: Lentiviral Particles

Product Name: TAB3 (NM_152787) Human Tagged ORF Clone Lentiviral Particle

Symbol: TAB3

Synonyms: MAP3K7IP3; NAP1

Mammalian Cell

Selection:

Puromycin

Vector: pLenti-C-Myc-DDK-P2A-Puro (PS100092)

NM 152787

Tag: Myc-DDK

ORF Size: 2136 bp

ORF Nucleotide

The ORF insert of this clone is exactly the same as(RC207646).

Sequence:

ACCN:

OTI Disclaimer: The molecular sequence of this clone aligns with the gene accession number as a point of

reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing

variants is recommended prior to use. More info

OTI Annotation: This clone was engineered to express the complete ORF with an expression tag. Expression

varies depending on the nature of the gene.

RefSeg: NM 152787.3

 RefSeq Size:
 6798 bp

 RefSeq ORF:
 2139 bp

 Locus ID:
 257397

 UniProt ID:
 08N5C8

Cytogenetics: Xp21.2

Domains: zf-RanBP, CUE

Protein Families: Druggable Genome





TAB3 (NM_152787) Human Tagged ORF Clone Lentiviral Particle - RC207646L3V

Protein Pathways: NOD-like receptor signaling pathway

MW: 78.7 kDa

Gene Summary: The product of this gene functions in the NF-kappaB signal transduction pathway. The

encoded protein, and the similar and functionally redundant protein MAP3K7IP2/TAB2, forms a ternary complex with the protein kinase MAP3K7/TAK1 and either TRAF2 or TRAF6 in

response to stimulation with the pro-inflammatory cytokines TNF or IL-1. Subsequent MAP3K7/TAK1 kinase activity triggers a signaling cascade leading to activation of the NF-

kappaB transcription factor. The human genome contains a related pseudogene.

Alternatively spliced transcript variants have been described, but their biological validity has

not been determined. [provided by RefSeq, Jul 2008]